# ADDITIONAL TRANSFERS OF CHINESE ACANTHOPANAX TO ELEUTHEROCOCCUS (ARALIACEAE)

# PING-SHENG HSU

Laboratory of Systematic and Evolutionary Botany
Institute of Botany
Chinese Academy of Sciences
Beijing 100093
PEOPLE'S REPUBLIC OF CHINA
SHENG-LI PAN

Faculty of Pharmacognosy, School of Pharmacy Shanghai Medical University Shanghai 200032 PEOPLE'S REPUBLIC OF CHINA

### **ABSTRACT**

The priority of *Eleutherococcus* Maxim. over *Acanthopanax* (Decne. & Planch.) Witte necessitates additional transfers of *Acanthopanax baoxinensis* X.P. Fang et C.K. Hsieh, *A. cissifolius* (Griff. ex Seem.) Harms var. *glaber* Y.R. Li, *A. nanpingensis* X.O. Fang et C.K. Hsieh, *A. giraldii* Harms var. *pilosulus* Rehd. and *A. yui* H.L. Li var. *villosus* Y.R. Li to the genus *Eleutherococcus*.

## RESUMEN

La prioridad de *Eleutherococcus* Maxim. sobre *Acanthopanax* (Decne. & Planch.) Witte necesita las transferencias adicionales al género *Eleutherococcus* de *Acanthopanax baoxinensis* X.P. Fang et C.K. Hsieh, *A. cissifolius* (Griff. ex Seem.) Harms var. *glaber* Y.R. Li, *A nanpingensis* X.P., Fang et C.K. Hsieh, *A. giraldii* Harms var *pilosulus* Rehd. y *A. yui* H.L. Li var *villosus* Y.R. Li.

Key words: Eleutherococcus, Acanthopanax, Araliaceae, new combinations

Acanthopanax was first used as a subgeneric name of Panax L. by Decaisne and Planchon (1854). It was later raised by Witte (1861, cf. Philipson 1979) and Miquel (1863) to generic rank. Acanthopanax (Decne. & Planch.) Witte and Eleutherococcus Maximowicz (1859) were recognized by Bentham & Hooker (1865), Forbes & Hemsley (1888), Nakai (1914), 1924, 1927), Poyarkova (1950), etc. as separate genera. However, Harms (1894, 1918) combined both genera and recognized Eleutherococcus in sectional rank under the genus Acanthopanax, a treatment that cannot be allowed under the current ICBN. Harms' treatment was followed by Rehder (1940, 1949), Li (1942), and many other floristic botanists in China, Japan and southeastern Asia (e.g. Philipson 1979). However, Stone (1980) transferred to Eleutherococcus from Acanthopanax the single Malayan

594 SIDA 15(4): 1993

species, *A. malayanus* Henderson, and affirmed the nomenclatural need for further changes of this kind to consolidate the name which has priority.

The late Hoo & Tseng (1978) accepted Harms' system (1918) (except sect. Kalopanax which was regarded as a distinct genus). They used Acanthopanax as the generic name, and noted that that name had been in common use for long periods and should be conserved over the earlier synonym Eleutherococcus. But they did not put forward such a proposal. Nearly all the floristic works as well as publications of pharmaceutical botany in China have followed Hoo & Tseng's treatment. However, Hu (1980) recognized *Eleutherococcus* to be the earlier validly published name for Acanthopanax, and transferred many East Asiatic taxa of Acanthopanax to Eleutherococcus. But she overlooked the publications of Hoo & Tseng (1965, 1978), and left quite a lot of taxa in Acanthopanax which were transferred by Ohashi in 1987 in a new and revised system of Eleutherococcus. Zhang and Liu (1988) published new combinations for four taxa of Chinese Eleutherococcus. But unfortunately three of their combinations were predated by Ohasi (1987). Since the generic name Acanthopanax has never been proposed to conserve against Eleutherococcus, transfers of taxa from that genus to the latter are necessary. In this paper, new combinations are published for those taxa still remaining in Acanthopanax in the Chinese flora.

- 1. Eleutherococcus baoxinensis (X.P. Fang et C.K. Hsieh) Hsu et S.L. Pan, comb. nov. *Acanthopanax baoxinensis* X.P. Fang et C.K. Hsieh, Bull. Bot. Res. 7(4): 89, fig. 2. 1987.
- 2. Eleutherococcus cissifolius (Griff. ex Seem.) Nakai var. glaber (Y.R. Li) Hsu et S.L. Pan, comb. nov. *Acanthopanax cissifolius* (Griff. ex Seem.) Harms var. glaber Y.R. Li, Acta Bot. Yunnan. 2(1):107. 1980.
- 3. Eleutherococcus nanpingensis (X.P. Fang et C.K. Hsieh) Hsu et S.L. Pan, comb. nov. *Acanthopanax nanpingensis* X.P. Fang et C.K. Hsieh, Bull Bot. Res. 7(4):87, fig. 1. 1987.
- 4. Eleutherococcus wilsonii (Harms) Nakai var. pilosulus (Rehd.) Hsu et S.L. Pan, comb. nov. Acanthopanax giraldii Harms var. pilosulus Rehd., J. Arnold Arbor. 9:99. 1928. E. giraldii (Harms) Nakai var. pilosulus (Rehd.) S.Y. Hu, J. Arnold Arbor. 61:109. 1980. A wilsonii Harms var. pilosulus (Rehd.) X.P. Fang et C.K. Hsieh, Bull. Bot. Res. 7(4):90. 1987.

Fang and Hsieh (1987) considered this variety belongs to *Acanthopanax* wilsonii Harms rather than to *A. giraldii* Harms. The present authors are in favor of their suggestion.

5. Eleutherococcus giraldii (Harms) Nakai var. villosus (Y.R. Li) Hsu et S.L. Pan, comb. nov. Acanthopanax yui H.L. Li var. villosus Y.R. Li, Fl Yunnan. 2:479. 1979. Shang (1985) considered Acanthopanax yui Li conspecific with A. giraldii Harms. The present combination is based on Shang's treatment.

### REFERENCES

- Bentham, G. and J.D. Hooker. 1865. Araliaceae, In Genera Plantarum 1:931-947.
- Decaisne, J. and J.E. Planchon. 1854. Esquisse d'une monographie des Araliacées. Revue Hort. IV. 3:104–109.
- FANG, X.P. and C.K. Hsieh. 1987. New taxa of *Acanthopanax* Miq. from Sichuan. Bull. Bot. Res. 7(4):87–92.
- FORBES, F.B. and W.B. Hemsley. 1888. Araliaceae, In Enum. Pl. China 1:337-343.
- HARMS, H. 1894. Araliaceae, In Nat. Pflanzenfam. III. 8:1-62 (Acanthopanax, 49-51).
- \_\_\_\_\_\_. 1918. Übersicht über die Arten der Gattung Acanthopanax. Mitt. Deutsch. Dendrol. Ges. 27:1–39, pls. 1–8.
- Hoo, G. and C.J. Tseng. 1965. Contributions to the Araliaceae of China. Acta Phytotax. Sin. Add. I. 129–176.
- \_\_\_\_\_. 1978. Araliaceae, In Flora Reipub. Pop. Sin. 54. Beijing.
- Hu, S.Y. 1980. Eleutherococcus vs. Acanthopanax. J. Arnold Arbor. 61:107-111.
- Li, H.L. 1942. Araliaceae of China. Sargentia 2:1-134.
- Maximowicz, C.J. 1859. Primitiae florae Amurensis. Mem. Acad. Sci. St.-Pétersb. Sav. Étr. 9:1–504, pls. 1–10, 1 map.
- Miquel, F.A.W. 1863. Araliaceae novae, adiecta aliarum specierum praesertim Indicarum revisione. Ann. Mus. Lugd.-Bat. 1:1–27.
- Nakai, T. 1914. Chosen shokubatsu 1. Seibido, Tokyo.
- . 1924. Araliaceae imperii Japonici. J. Arnold Arbor. 5:1–36.
- Oнashi, H. 1987. *Eleutherococcus* (Araliaceae) a new system and new combinations. J. Jap. Bot. 62(12):353–361.
- \_\_\_\_\_\_. 1989. Eleutherococcus (Araliacaeae) of Japan. J. Jap. Bot. 64(7):216–218.
- Philipson, W.R. 1979. Araliaceae I, In Flora Malesiana I. 9:1–105. Martinus Nijhoff, Hague, Netherlands.
- Poyarkova, A.I. 1950. Araliaceae, In Fl. SSSR 16. Moskva.
- REHDER, A. 1940. Manual of cultivated trees and shrubs. Ed. 2. MacMillan, New York.
- \_\_\_\_\_\_. 1949. Bibliography of cultivated trees and shrubs. Arnold Arboretum, Jamaica Plain, Massachusetts.
- Shang, C.B. 1985. New taxa and some revisions about the Araliaceae of China. J. Nanjing Inst. Forest. 2:15–28.
- Stone, B.C. 1980. A nomenclatural change in Malaysian Araliaceae: *Eleutherococcus malayanus* (Henderson) comb. nov. instead of *Acanthopanax malayanus* Henderson. Malaysian Forester 43(3):395.
- ZHANG, K.L. and H.X. Liu. 1988. Studies on the genus *Eleutherococcus* Maxim. of Gansu. J. North Normal Coll. (Nat. Sci. Ed.) Suppl. 1:10–16.